GBiot-FHO cost-effectively provides a high accuracy of measurement close to the continnuous monitoring official method) implemented by the MoE and local governments in Japan

and the second se				
Power Requirements	100/200VAC (for overseas use) 12VDC (for solar operation)			
Power Consumption	approx. 5W (10W at peak)			
Data Communication	Wi-Fi (LPWA as an option)			
Measurement Range /SO2	0 ~ 1000ppb			
Measurement Range /CO	0 ~ 20 ppm			
Measurement Range /03	$0{\sim}500{ m ppb}$			
Measurement Range / PM	$0{\sim}500ug/m^3$			
Measurement Range /NO2	$0{\sim}250\text{ppb}$			
Weather Sensor	Temp. Hum. Press. WD. WS. Rainfall Solar Radiation UV int.			
Sensor Life	$1\!\sim\!5$ years (depend on sensors and frequency of use)			
X Specifications of device and service are subject to change without notice				







% Specifications of device and service are subject to change without notice




GREEN BLUE

Contact

https://gbiot.jp/ https://greenblue.co.jp/ Phone 81-45-322-3155 e-mail sales@greenblue.co.jp


Comparison in PM25




Comparison results between GBiot and a light scattering dust analyzer with the same measuring principle are shown below.


As can be seen from the scatter plots and regression equations, GBiot and DustTrak showed a high correlation of $R^2 = 0.98$ or higher.

※ DustTrak is a type of light scattering dust analyzer




Comparison in O3





Comparison tests between GBiot and the official method for O3 were conducted this winter and summer at Green Blue's Tokyo office and surrounding monitoring stations.

As shown in the scatter plots and regression equations, the comparison between GBiot and the official method showed a high correlation of $R^2 = 0.9031$.



Installation and Maintenance

	GDIO	t-FH0 Dashboar		1022-07-10-07:40:23				
Dutbert	Each location name is hidden							
. Clinet	-		-	-sta	-		Are -	-
-A Chief	6-19	-944.00	349	100.00	1418	441414	101	Junit of
Desettad -	114	hile	114	10.00.00	14	11-13-	ive	A7.0
	Linguage	of sectors.	LATTING	14.795.019	Lawran	ILINOIDA .	Lansage .	AL PROPERTY.
12 hour	- unitial	10.05701	and at	A.4570	and the	8.040	chanted	and regard
	80	Ange 41.0	401	ton upon	4462.5	1.41 -0091	4403	J.M. opped
		1 apr	14	10.00	100	10.00	10	1.00
	41	21.000	687	3.00	100	1.000	40	1.00
	terminate and	41.5	THE REAL PROPERTY AND	9.9 K	mental	21.64	(BARRATUR)	44.0
	Autom.	40.76	- March	AL 4.	and the second s	142 m		de 16.
	-	8.401	-	199. pt	-	-57. 44		64.100
		are an		10 M		107.00		10.00

- Remote data check is possible at any point where online communication environment is organized.
- With GBiot's software, data can be easily graphed and downloaded.
- Video manuals are available to support the installation and maintenance of GBiot.





Considerations

Classification	Things to be considered in system construction and operation
Installation Site	 Representative point in the area, avoiding the influence of specific sources, air currents, and direct sunlight Ensuring stability of equipment (e.g., fixation by anchors)
Power Supply	 Securing a stable AC power supply (including measures against lightning strikes)
Data Communication	 Ensuring a stable communication environment (e.g., Wi-Fi)
Security	 Prevension against theft of equipment and data
Maintenance	 Securing regular maintenance personnel and spare parts (frequency of work: once per 6 months to 1 year) Emergency maintenance personnel and parts availability (irregular)
Calibration	 Calibration should be performed by running the GBiot in parallel with the reference instrument of the official method




Observation of air pollution by GBiot in Myanmar

Air pollutants such as PM2.5 and NO2 have been observed by GBiot in 4 cities in Myanmar by a certain agency.
 Among them, in Yangon City, GBiot and two sets of official methods are being operated in parallel.



Practice for data collection and visualization of air pollution in Vietnam

With the local counterparts, Green Blue collected data of air pollution and video around a lake in Vietnam using three monitoring stations and two motorcycles from September to November 2020. Based on data analysis and visualization, the situation of air pollution around the lake was grasped. Note: Based on demonstration experiments of NICT xData platform



All the sensorboxes for air pollution used in the demonstration experiment are made by Green Blue

